

Searching New Physics in Muonium Atoms

Klaus P. Jungmann¹

Physikalisches Institut, Universität Heidelberg, Philosophenweg 12, D-69120 Heidelberg, Germany

Abstract. Quantum Electrodynamics describes electromagnetic interactions of free muons and of muon and electron in the muonium atom to very high accuracy. Contributions of other and yet unknown interactions can be investigated in precision experiments, e.g. by searching for muonium-antimuonium conversion and by measuring the muon magnetic anomaly.

1. Muonium-antimuonium conversion

The hydrogenlike muonium atom ($M = \mu^+ e^-$) consists of two leptons from different generations. The close confinement of the bound state offers excellent opportunities to explore precisely fundamental electron-muon interaction. The dominant part of the binding in this atom is electromagnetic which can be calculated to very high accuracy in the framework of quantum electrodynamics (QED). Indeed, precision experiments of electromagnetic transitions in this system have been employed both for testing the validity of bound state QED calculations and for determining most accurate values of fundamental constants [1]. The present 36 ppb precision of measurements of the hyperfine structure of muonium is already a factor of two below the 50 ppb contribution of strong interactions arising from hadronic vacuum polarization. New experimental results on this transition from recently completed measurements at the Los Alamos Meson Physics Facility (LAMPF) are expected to reveal a 15 ppb contribution of weak interaction due to Z_0 boson exchange.

Since the effects of all known fundamental forces in this system are calculable very well, it renders the possibility to search sensitively for yet

¹ E-mail: jungmann@physi.uni-heidelberg.de





Figure 1. Muonium-antimuonium conversion in theories beyond the standard model. The interaction could be mediated by (a) a doubly charged Higgs boson Δ^{++} [6], (b) heavy Majorana neutrinos [5], (c) a neutral scalar Φ_N [8], e.g. a supersymmetric τ -sneutrino $\tilde{\nu}_\tau$ [7], or (d) a dileptonic gauge boson X^{++} [9].

unknown interactions between both particles. A conversion of muonium into its antiatom ($\bar{M} = \mu^- e^+$) would violate additive lepton family number conservation. Although no symmetry behind this experimentally established law has been found, the process is not provided in standard theory. However, muonium-antimuonium conversion appears to be natural in many speculative theories (see Fig. 1), which try to extend the standard model in order to explain some of its yet not well understood features like parity violation in weak interaction and particle mass spectra. An experiment has been set up to search for spontaneous muonium-antimuonium conversion at the Paul Scherrer Institute (PSI) in Villigen, Switzerland (Fig. 1) [2]. It uses the powerful signature developed in an experiment at LAMPF, which requires the coincident identification of both particles forming the antiatom in its decay [3, 4].

Muonium atoms were produced by stopping a beam of "surface" muons in a SiO_2 powder target, where a 60 % fraction of them forms muonium atoms by electron capture some 5% of which diffuse through the target surface with thermal energies. Energetic electrons from the decay of the μ^- in the antiatom can be observed in a magnetic spectrometer at 0.1 T magnetic field consisting of five concentric multiwire proportional chambers and a 64 fold segmented hodoscope. The positron in the atomic shell of the antiatom is left behind after the decay with 13.5 eV average kinetic energy. It can be accelerated to 8 keV in a two stage electrostatic device and guided in a magnetic transport system onto a position sensitive microchannel plate detector (MCP). Annihilation radiation can be observed in a 12 fold segmented pure CsI calorimeter surrounding the MCP.

The relevant measurements were performed during in total 6 month

Figure 2. Top view of the apparatus to search for muonium-anti-muonium conversion at PSI [2].



distributed over 4 years. The muonium production was monitored regularly by reversing all electric and magnetic fields of the instrument every five hours for a duration of 20 minutes. Targets had to be replaced twice a week because of an observed deterioration of muonium production on a time scale of about one week. In the course of the experiment $5.7 \cdot 10^{10}$ muonium atoms were observed in the interaction volume for antimuonium decays. There was one event which passed all required criteria, i.e. fell within a 99% confidence interval of all relevant distributions (Fig. 1). The expected background due to accidental coincidences is 1.7(2) events. The preliminary




Figure 3. Time of flight (TOF) and vertex quality for a muonium measurement (left) and all data of 1996 while searching for antimuonium in 1996 (right). One event falls into the indicated 3 standard deviations area.

combination of all analyzed data results in an upper limit for the conversion

probability in a 0.1 T magnetic field of $P_{\overline{MM}} \leq 8 \cdot 10^{-11}$. For an assumed effective (V-A)x(V-A) type four fermion interaction this corresponds to an upper limit for the coupling constant of $G_{\overline{MM}} \leq 3 \cdot 10^{-3} G_F$, where G_F , (90 % C.L.) is the weak interaction Fermi coupling constant [10]. This includes a correction for the magnetic field [11].

This new result allows to rule out definitively a certain Z_8 model with more than 4 generations of particles [8] and to set a new lower limit of 2.6 TeV/c² on the masses of dileptonic gauge bosons in GUT models which is well beyond the value extracted from high energy Bhabha scattering [9]. It can be further shown in the framework of minimal left right symmetric and supersymmetric models that lepton number violating muon decay ($\mu^+ \rightarrow e^+ + \nu_\mu + \overline{\nu}_e$) is not an option for explaining the excess neutrino counts in the LSND neutrino experiment at Los Alamos [12].

2. Magnetic anomaly of the muon

The muon magnetic anomaly, i.e. the deviation of the muon's g-factor from the Dirac value 2, has been measured in a series of three experiments at CERN [13] to an accuracy of 7 ppm. Although this quantity is determined, as in the case of the electron, mostly by photon and electron-positron fields, the effects of heavier particles is enhanced compared to the electron cases by the square of the mass ratio $(m_\mu/m_e)^2 \approx 4 \cdot 10^4$. The contribution of the strong interaction is 60 ppm and the weak contribution is expected to be 1.3 ppm.

This summer a new experiment [14] was successfully brought into operation at the Brookhaven National Laboratory (BNL) in Upton, USA, the ultimate goal of which is a determination of the muon magnetic anomaly to 0.35 ppm. At this level of sensitivity numerous theories beyond the

Table 1. Sensitivity to new physics of the new g-2 experiment at BNL.

New Physics	Sensitivity			Other Experiments
Muon substructure	Λ	$\geq$	5 TeV	LHC similar
excited muon	m_{μ^*}	$\geq$	400 GeV	LEP II similar
$W^\pm$ -boson substructure	Λ	$\geq$	400 GeV	LEP II $\sim$ 100-200 GeV
$W^\pm$ anomalous magnetic moment	a_W	$\geq$	0.02	LEP II $\sim$ 0.05, LHC $\sim$ 0.2
Supersymmetry	$m_{\widetilde{W}}$	$\leq$	130 GeV	Fermilab $p\overline{p}$ similar
right handed $W_R^\pm$ -bosons	$m_{W'}$	$\leq$	250 GeV	Fermilab $p\overline{p}$ similar
Muon el. dipole moment	D_μ	$\leq$	$4 \cdot 10^{-20} ecm$	

^a for substructure $\Delta a_\mu \sim m_\mu^2/\Lambda^2$

standard model can be tested, respectively their parameters can be significantly restricted (see Table 1) [14, 15]. The potential of this high precision experiment reaches in part the one of high energy physics experiments and exceeds it even some cases.

3. Conclusions and Acknowledgments

Precision experiments on muons and muonium can contribute to probe sensitively physics beyond the standard model in a complementary way to high energy physics approaches. Maybe ultimately they can contribute to resolve the question of the difference in nature between muon and electron.

This work was supported by the German Bundesminister für Bildung und Forschung and by a NATO research grant.

References

- [1] Hughes V W and zu Putlitz G 1990 in: *Quantum Electrodynamics* (Singapore: World Scientific, Kinoshita T, ed.) 822; Jungmann K 1994 in: *Atomic Physics 14* (New York: AIP Press, Wineland D et al.,ed.) 102
- [2] Abela R et al. 1996 *Phys. Rev. Lett.* **77** 1951
- [3] Matthias B et al. 1991 *Phys. Rev. Lett.* **66** 2716
- [4] Willmann L and Jungmann K 1997 (in print) in: *Atomic Physics Methods in Modern Research* (Heidelberg: Springer)
- [5] Halprin A 1982 *Phys. Rev. Lett.* **48** 1313
- [6] Herczeg P and Mohapatra R N 1992 *Phys. Rev. Lett.* **69** 2475
- [7] Mohapatra R N 1992 *Z. Phys. C* **56** 117; Halprin A and Masiero A 1993 *Phys. Rev. D* **48** 2987
- [8] Hou W S and Wong G G 1996 *Phys. Rev. D* **53** 1537
- [9] Fujii H et al. 1994 *Phys. Rev. D* **53** 560
- [10] Meyer V et al. 1997 in: Proceedings of the International Conference on Intersections between Particle and Nuclear Physics, Big Sky
- [11] Wong G G and Hou W S 1995 *Phys. Lett. B* **357** 145; Horikawa K and Sasaki K 1996 *Phys. Rev.* **53** 560
- [12] Herczeg P this conference and priv. com.; Louis W this conference
- [13] Farley F J M *Z. Phys. C* **56** 88
- [14] Roberts B L 1997 loc.cit [4]; Miller J et al 1997 loc. cit. [10]
- [15] Kinoshita T and Marciano W J loc.cit. [1] p. 419; Roberts B L 1992 *Z. Phys. C* **56** 101